



## *Aspergillus homomorphus*, a first global record from millet grains

Nemmat Hussein<sup>1</sup>, Abdel-Hafez SII<sup>1</sup>, Abdel-Sater MA<sup>1</sup>, Ismail MA<sup>1</sup> and Eshraq AL-Amrey<sup>1,2</sup>

<sup>1</sup>Botany & Microbiology Department, Faculty of Science, Assiut University, P.O. 71516, Assiut, Egypt.

<sup>2</sup>Microbiology Department, Faculty of Applied Science, Taiz University, Yemen.

Hussein N., Abdel-Hafez SII, Abdel-Sater MA, Ismail MA, AL-Amrey E. 2017 – *Aspergillus homomorphus*, a first global record from millet grains Current Research in Environmental & Applied Mycology 7(2), 82–89, Doi 10.5943/cream/7/2/4

### Abstract

During surveys of mycobiota associated with millet (*Pennisetum glaucum* L.) grains from Taiz Governorate, Yemen, an interesting isolate of *Aspergillus* section *Nigri* was isolated. It was recorded once on Czapek's agar at 28°C. The isolate was examined and identified macroscopically, microscopically and genotypically as *Aspergillus homomorphus*. It is recorded for the first time from millet grains and for only the third time since its original description in 1994. The strain was deposited at Assiut University Mycological Centre (AUMC), and the ITS sequence was deposited in the National Centre of Biotechnology Information (GenBank).

**Key words** – Assiut University Mycological Centre – Czapek's agar – DNA sequencing – Section *Nigri* – Yemen.

### Introduction

Members of *Aspergillus* section *Nigri* (black aspergilli) are important in biotechnological processes as well as in biodeterioration. Some strains of black aspergilli are often misidentified as *A. niger* due to the difficulties of identifying the species in this group (Samson et al. 2007). Recent surveys on the mycobiota of distinct commodities revealed new species of section *Nigri* unknown to science (Perrone et al. 2007). For many years black aspergilli, and indeed all *Aspergillus* spp. were identified to species by morphological criteria (Raper & Fennell 1965, Pitt & Hocking 2009, Oliveri & Catara 2011). such as colour, density and colony appearance and microscopic observations (conidial head, conidiophore and conidial characters). The taxonomy of the section was revised by Abarca et al. (2004). and 15 taxa were accepted in the last critical revisions: *A. aculeatus*, *A. brasiliensis*, *A. carbonarius*, *A. costaricaensis*, *A. ellipticus*, *A. japonicus*, *A. foetidus*, *A. heteromorphus*, *A. homomorphus*, *A. lacticoffeatus*, *A. niger*, *A. piperis*, *A. sclerotioniger*, *A. tubingenensis* and *A. vadensis* (Samson et al. 2004, 2007). Subsequently, many species of this section were described based mainly on their genotypic characteristics (Serra et al. 2006, Noonim et al. 2008, Perrone et al. 2008, Sørensen et al. 2011, Varga et al. 2011, Jurjević et al. 2012). *A. homomorphus* was not only separated from related species in the section by morphological distinction but also by molecular analyses (Kallow et al. 2006). Among morphological differences, this species has conidia that are 5–7 µm in diameter, which separates it from *A. carbonarius* (7–9 µm diam.) and *A. niger* aggregate (3–5 µm diam.) (Samson & Varga 2007). Perrone et al. (2007).

reported that nine different black *Aspergillus* species are often difficult to identify with classical methods, but can be distinguished by molecular techniques.

During mycobiotic surveys carried out continuously in the Mycological Laboratory at the Department of Botany and Microbiology, Faculty of Science, Assiut University, an unusual black strain of *Aspergillus* was isolated from millet grains collected from Taiz City, Yemen and placed on Czapek's agar medium at 28°C. It was identified based on macro – and micromorphology and genotypically as belonging to *Aspergillus homomorphus*. This species was first isolated and described (invalidly) together with the closely related species (*A. pseudo-heteromorphus*), as new species from soil samples collected near the Dead Sea (Steiman et al. 1994). Samson et al. (2004). validated *A. homomorphus* and synonymized *A. pseudo-heteromorphus* with *A. homomorphus* since they were identical in both extrolites and DNA sequences. *A. homomorphus* was later recorded from a soil sample in Mexico (Borrego–Terrazas et al. 2014). and soil in the arid regions of Oman (Al-Sadi /ncbi website).

## **Materials & Methods**

### **Source of the isolate**

The isolate was obtained during surveys of grain-borne fungi of millet (*Pennisetum glaucum* L.) samples collected from Taiz Governorate, Yemen. For isolation of fungi the grains were placed on Czapek's agar medium and incubated at 28°C.

### **Media and growth conditions for identification**

The isolate was cultivated and examined on the following standard media: Czapek's yeast autolysate agar (Samson & Pitt 1985). Czapek's agar (Raper & Thom 1949). Czapek's with 20% sucrose (Cz20S), (Raper & Fennell 1965). malt extract agar, (MEA) ( Blakeslee 1915). malt yeast with 40% sucrose agar, (M40Y), (Raper & Fennell 1965). glycerol 25% nitrate agar, (G25N), (Pitt 1973). creatine sucrose agar, (CREA), (Frisvad 1985). tannin sucrose agar, (TAN), (Thrane 1986). urea agar (UA), mannitol agar, (MA), (Brayford & Bridge 1989). and Czapek's starch agar (CzSA) in which sucrose was replaced by soluble starch (Bridge 1985). Three replicates of 3-pointed inoculated plates of all media were incubated at 25°C except for CYA, plates were incubated at 5°, 25°, 37° and 45°C. Growth rates were recorded on CYA, Cz and MEA after 7 days of incubation. Assessment of growth on media with reduced water activity (Cz20S, G25N and M40Y). was also carried out after 7 days?. Turning the UA medium pink was referred as urease positive. A positive amylase result was indicated by the appearance of a clear zone around the colony after flooding with iodine solution (Cowan 1974).

### **Growth rates**

Colony diameters were measured on all media and all tested temperatures. The results on MA were assessed by growth and acid production turning phenol red pH indicator from red to yellow. Growth and base production on CREA were also recorded by visible colour change of medium from purple to yellow.

### **Phenotypic characters**

Assessment of colony characteristics (colony texture, colour, exudate, reverse and soluble pigment) was carried out. The Methuen Handbook of Colour (Kornerup & Wanscher 1989). was used for evaluation of colour. Microscopic measurements of conidial heads, stipes (conidiophores), vesicles, supporting cells (metulae), conidiogenous cells (phialides) and conidia were determined. Surface and colour of the stipe and conidia were also recorded.

### **Sample preparation for Scanning Electron topography Microscope (SEM)**

Discs from culture grown on CYA at 25°C for 5 days were soaked in a fixative solution at pH 7.2. The discs were then washed in cacodilate buffer, post-fixed in 1% osmium tetroxide solution and water for 1 hour, followed by dehydration in increasingly more concentrated acetone solutions. Afterwards, the samples were dried in a desiccator containing silica. The specimens were clustered on a film of aluminum foil, then coated with gold in a sputter and observed in a scanning electron microscope, SEM (JEOL, JSM-5400 LV unit, Assiut University, Assiut, Egypt) (Silva et al. 2011).

### Photography

Growth on all agar media as well as microscopic structures were photographed. Moreover, a number of images from SEM were digitally produced and registered at variable magnifications.

### Genetic identification

The isolate was grown on MEA plates and incubated at 28°C for 4 days. Mycelial growth was scraped off and immersed in 100 µl of distilled water and boiled at 100°C for 15 minutes. It was then sent to SolGent Company, Daejeon, South Korea for DNA extraction and sequencing. Internal transcribed spacer (ITS) sequences of nuclear rDNA were amplified using universal primers ITS1 (5' – TCC GTA GGT GAA CCT GCG G – 3') and ITS4 (5'– TCC TCC GCT TAT TGA TAT GC –3'). Then amplification was carried out using the polymerase chain reaction, PCR (ABI, 9700). The PCR reaction mixtures were prepared using SolGent EF-Taq as follows: 10X EF-Taq buffer 2.5 µl, 10 mM dNTP (T) 0.5 µl, primer (F-10p) 1.0 µl, primer (R-10p) 1.0 µl, EF-Taq (2.5U) 0.25 µl, template 1.0 µl, DW to 25 µl. Then the amplification was conducted using the following PCR reaction conditions: one round of amplification consisting of denaturation at 95°C for 15 min followed by 30 cycles of denaturation at 95°C for 20 sec, annealing at 50°C for 40 sec and extension at 72°C for 1 min, with a final extension step of 72°C for 5 min. The PCR products were then purified with the SolGent PCR purification Kit-Ultra (SolGent, Daejeon, South Korea) prior to sequencing. The purified PCR products were reconfirmed (using size marker) by electrophoresis of the PCR products on 1% agarose gel. The bands were eluted and sequenced. Each sample was sequenced in the forward and reverse direction. The sequence alignments were performed using Clustal W analysis and the phylogenetic analysis of ITS retrieved from GenBank database (Moubasher et al. 2016). Contigs were created from the sequence data using CLCBio Main Workbench program. The sequence obtained from the target isolate was further analyzed using BLAST from the National Center of Biotechnology Information (NCBI) website. The sequence was subjected to Clustal W analysis using MegAlign (DNASstar) software version 5.05 for the phylogenetic analysis. The ITS sequence was deposited in Genbank (Thompson et al. 1994).

### Results

#### *Colony characteristics of Aspergillus homomorphus Steiman, Guiraud, Sage & Seigle-Murandi ex Samson and Frisvad*

**On Cz at 25°C after 7 days:** Colonies 20–28 mm diam, floccose, margin white, conidiogenesis moderate to heavy; colour black (M. 6E–F1) to carbon black (Kornerup & Wanscher 1989). sclerotia absent, exudate, reverse and soluble pigment absent.

**On CYA at 25°C after 7 days:** Colonies 45–64 mm diam, floccose, margin white, conidiogenesis as on Cz; colour carbon black; sclerotia absent; exudate and soluble pigment absent; reverse colour yellowish white, pale yellow to pastel yellow (M. 2A2–4).

**On CYA at 5° and 45°C after 7 days:** No growth.


**On CYA at 37°C after 7 days:** Colonies 30–50 mm diam, colony texture, colour, conidiogenesis and reverse as on CYA at 25°C after 7 days.

**On MEA at 25°C after 7 days:** Colonies 40–60 mm diam, floccose, margin white, conidiogenesis moderate; colour black; sclerotia; exudate, reverse and soluble pigment absent (Fig 1). Colony diameters of the strain on other media was presented in (Table 1).

**Table 1** Growth rate of *Aspergillus homomorphus* on different media.

Medium	Colony diameter (in mm)
Malt yeast with 40% sucrose agar (M40Y)	40 – 55
Czapek´s with 20% sucrose (Cz20S)	32 – 38
Creatine sucrose agar (CREA)	21 – 32
Tannin sucrose agar (TAN)	24 – 35
Glycerol 25% nitrate agar (G25N)	2 – 10
Czapek´s starch agar (CzSA)	22-24
Urea agar (UA)	35 – 48
Mannitol agar (MA)	17 – 25

Note: The fungus showed base production on CREA, amylase production, but negative urease activity. In respect to MA, acid production was not observed.




**Fig 1** – *Aspergillus homomorphus* AUMC 10521 grown on CYA at 5°, 25°, 37° and 45°C; and MEA, Cz, Cz20%S, G25N, mannitol, creatine, tannin, M40Y and urea at 25°C.

### *Microscopic features*


On CYA, conidial heads biseriate, radiate to globose, 80–180 µm diam, splitting with age into defined columns, stipes smooth-walled, light yellowish brown colour below the vesicle, 400–875 x 9–15 µm, sometimes with septum below the vesicle, vesicle globose, subglobose to less commonly spathulate, brown, 27–50 µm diam, but slightly smaller in uniseriate heads (20–45 µm diam), metulae present, 7–8 µm long; phialides 9–15 x 4–5 µm, conidia globose to subglobose,



conspicuously spiny to echinate at maturity, mature conidia dark brown, 5–7  $\mu\text{m}$  diam, but up to 8  $\mu\text{m}$  diam on MEA, with spines up to 2  $\mu\text{m}$  in length (Figs 2 – 11).



**Figs 2–7** – Microscopic structure of *A. homomorphus* AUMC 10521. 2–4 Conidial heads. 5–7 Conidia.




**Figs 8–11** – Scanning electron micrographs of *A. homomorphus* AUMC 10521. 8 Stipes and conidial head. 9 Phialides and conidia. 10–11 Conidia.

### ***Molecular identification***

ITS sequence of *A. homomorphus* AUMC 10521 showed high similarity (> 99%) with GenBank accession numbers NR-077189 = CBS 101889<sup>T</sup> (type strain), EU821330 = CBS 101889<sup>T</sup>, KU94906 = SQU-MA06 and KJ888832 = TAA50 and 94.2–94.7 with *A. aculeatinus* (NR\_135417 = CBS 121060<sup>T</sup>), *A. indologenus* (AJ28305 = CBS 114.80), *A. uvarum* (NR\_135330 = CBS 121591<sup>T</sup>), *A. brunneoviolaceus* (NR\_138279 = NRRL4912<sup>T</sup>), *A. assiutensis* (JN393254 = AUMC

5748<sup>T</sup>), *A. japonicus* (NR\_131268 = CBS 114.51<sup>T</sup>) and *A. aculeatus* (NR\_111412 = CBS 172.66<sup>T</sup>) (Fig 12).



**Fig 12** – Phylogenetic tree based on DNA sequence data of ITS region of *Aspergillus homomorphus* AUMC 10521, compared with 13 reference strains in GenBank of closely related species belonging to *Aspergillus* section *Nigri* and *A. candidus* as outgroup.

*A. homomorphus* is most similar phenotypically to *A. ibericus* and *A. carbonarius*; the three species have black biseriate aspergilli with long stipes and relatively larger conidia (> 5 µm) when compared to species of the *A. niger* aggregate (conidia 3–5 µm diam), and smaller when compared to *A. carbonarius* (7–10 µm) (Kallow et al. 2006, Perrone et al. 2007). Samson et al. (2007). studied morphological characteristics of the type strains of *Aspergillus* Section *Nigri* and reported that several species produce large conidia up to 7–9 µm including *A. carbonarius*, *A. ibericus*, *A. sclerotioniger*, *A. homomorphus* and *A. scleroticarbonarius*, but *A. homomorphus* and *A. ibericus* produce conidia with spiny appearance. Other characters may help in the distinction of these species such as the dark or olive green colony reverse on Cz of *A. carbonarius* (Raper & Fennell 1965), versus absence of colour in the reverse in *A. homomorphus* and *A. ibericus*. Furthermore, on MEA, the latter two species grow faster than *A. carbonarius* (Raper & Fennell 1965). *A. ibericus* produces sclerotia whereas *A. homomorphus* does not (Frisvad et al. 2014). In agreement with the current results, *A. homomorphus* could grow at 36°C, but not at 45°C (Meijer et al. 2011).

CREA medium can be used as a semi-selective medium for classification of black aspergilli into groups (Samson et al. 2007). The present isolate grew and produced acid from CREA (turning the medium from purple to yellow). *A. homomorphus* showed colony diameter less than 40 mm at 37°C, short metulae, echinate conidia (spines up to 1.5 µm).

Blast search revealed a similarity of the current strain of *A. homomorphus* with the uniseriate aspergilli; *A. aculeatinus*, *A. aculeatus*, *A. assiutensis*, *A. brunneoviolaceus*, *A. indologenus*, *A. japonicus* and *A. uvarum* ranging from 94.2–94.7%. In similar studies, *A. homomorphus* and uniseriate species were located in either one subclade or two neighboring clades (Samson et al. 2004, Silva et al. 2011).

AUMC 10521 strain exhibited slightly smaller vesicles (up to 50 µm) than its original description of Steiman et al. (1994) (up to 65 µm). The current strain of *A. homomorphus* was isolated from millet grain and thus, the species is not limited to soil (Steiman et al. 1994, Borrego–Terrazas et al. 2014, Al–Sadi /ncbi website)

## References

- Abarca ML, Accensi F, Cano J, Cabañes FJ. 2004 – Taxonomy and significance of black aspergilli. *Antonie van Leeuwenhoek Int. Journal of Genetic Molecular Microbiology* 86(1), 33–49.
- Blakeslee A. 1915 – Lindner's roll tube method of separation cultures. *Phytopathology* 5, 68–69.
- Borrego Terrazas JA, Lara Victoriano F, Flores Gallegos AC, Veana F et al. 2014 – Nucleotide and amino acid variations of tannase gene from different *Aspergillus* strains. *Canadian Journal of Microbiology* 60(8), 509–516.
- Brayford D, Bridge PD. 1989 – Differentiation of *Fusarium oxysporum* from *Fusarium solani* by growth and pigmentation on media containing sugar alcohols. *Letters in Applied Microbiology* 9(1), 9–12.
- Bridge PD. 1985 – An evaluation of some physiological and biochemical methods as an aid to the characterization of species of *Penicillium* subsection Fasciculata. *Microbiology* 131(8), 1887–1895.
- Cowan ST. 1974 – Cowan and Steel's Manual for the Identification of medical bacteria, 2<sup>nd</sup> edn. Cambridge University Press, Cambridge, pp. 250.
- Frisvad JF. 1985 – Creatine sucrose agar, a differential medium for mycotoxin producing terverticillate *Penicillium* species. *Letters in Applied Microbiology* 1, 109–113.
- Frisvad JC, Petersen LM, Lyhne EK, Larsen TO. 2014 – Formation of sclerotia and production of indoloterpenes by *Aspergillus niger* and other species in section Nigri. *Plos One*, 9(4), 1–11.
- Jurjević Z, Peterson SW, Stea G, Solfrizzo M et al. 2012 – Two novel species of *Aspergillus* section Nigri from indoor air. *IMA Fungus* 3(2), 159–173.
- Kallow W, Santos I, Erhard M, Serra R et al. 2006 – *Aspergillus ibericus*: A new Species of Section *Nigri* Characterised by MALDI-TOF MS. *International Mycological Congress* 8, 189–193.
- Kornerup A, Wanscher JH. 1989 – Methuen Handbook of Colour. 3<sup>rd</sup> edition, London, pp. 252.
- Meijer M, Houbraken JAMP, Dalhuijsen S, Samson RA et al. 2011 – Growth and hydrolase profiles can be used as characteristics to distinguish *Aspergillus niger* and other black aspergilli. *Studies in Mycology* 69, 19–30.
- Moubasher AH, Abdel Sater MA, Soliman Z. 2016 – Biodiversity and molecular characterization of yeast and filamentous fungi in the air of citrus and grapevine plantations in Assiut area, Egypt. *Mycosphere* 7(3), 236–261.
- Noonim P, Mahakarnchanakul W, Varga J, Frisvad JC et al. 2008 – Two novel species of *Aspergillus* section *Nigri* from Thai coffee beans. *International Journal of Systematic and Evolutionary Microbiology* 58, 1727–1734.
- Oliveri C, Catara V. 2011 – Mycoflora and Biodiversity of Black Aspergilli in Vineyard Eco-Systems. *The Dynamical Processes of Biodiversity- Case Studies of Evolution and Spatial Distribution*, pp. 259.
- Perrone G, Susca A, Cozzi G, Ehrlich K et al. 2007 – Biodiversity of *Aspergillus* species in some important agricultural products. *Studies in Mycology* 59, 53–66.
- Perrone G, Varga J, Susca A, Frisvad JC et al. 2008 – *Aspergillus uvarum* sp. nov., an uniseriate black *Aspergillus* species isolated from grapes in Europe. *International Journal of Systematic and Evolutionary Microbiology* 58, 1032–1039.
- Pitt JI. 1973 – An appraisal of identification methods for *Penicillium* species: novel taxonomic criteria based on temperature and water relations. *Mycologia* 65, 1135–1157.
- Pitt JI, Hocking AD. 2009 – *Fungi and Food Spoilage*, 3<sup>rd</sup> ed. Springer- Verlag New York Inc. pp. 540.
- Raper KB, Fennell DI. 1965 – The genus *Aspergillus*. Baltimore, Williams and Wilkins, pp. 686.
- Raper KB, Thom C. 1949 – A manual of *Penicillium*. Williams and Wilkins, Baltimore, USA.
- Samson RA, Pitt JI. 1985 – *Advances in Penicillium and Aspergillus systematics*, Plenum Press, London, pp. 455–460.

- Samson RA, Varga J. 2007 – *Aspergillus* systematic in the genomic era. *Studies in Mycology* 59, 1–206.
- Samson RA, Houbraken J, Kuijpers A, Frank JM et al. 2004 – New ochratoxin A or sclerotium producing species in *Aspergillus* section *Nigri*. *Studies in Mycology* 50, 45–61.
- Samson RA, Noonim P, Meijer M, Houbraken J et al. 2007 – Diagnostic tools to identify black aspergilli. *Studies in Mycology* 59, 129–145.
- Serra R, Cabanes FJ, Perrone G, Castella G et al. 2006 – *Aspergillus ibericus*: a new species of section *Nigri* isolated from grapes. *Mycologia* 98 (2), 295–306.
- Silva DM, Batista LR, Rezende EF, Fungaro MHP et al. 2011 – Identification of fungi of the genus *Aspergillus* section *Nigri* using polyphasic taxonomy. *Brazilian Journal of Microbiology* 42(2), 761–773.
- Sørensen A, Lubeck PS, Lubeck M, Nielsen KF et al. 2011 – *Aspergillus saccharolyticus* sp. nov., a black *Aspergillus* species isolated in Denmark. *International Journal of Systematic and Evolutionary Microbiology* 61, 3077–3083.
- Steiman R, Guiraud P, Sage L, Seigle Murandi F. 1994 – New strains from Israel in the *Aspergillus niger* group. *Systematic and applied microbiology* 17(4), 620–624.
- Thompson JD, Higgins DG, Gibson TJ. 1994 – CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic acids research* 22(22), 4673–4680.
- Thrane U. 1986 – The ability of common *Fusarium* species to grow on tannin-sucrose agar. *Letters in Applied Microbiology* 2, 33–36.
- Varga J, Frisvad JC, Kocsubé S, Brankovics B et al. 2011 – New and revisited species in *Aspergillus* section *Nigri*. *Studies in Mycology* 69, 1–17.